Geoscience Canada



The Rice Lake Mine Trend, Manitoba:

Regional Setting, Host-rock Stratigraphy, and Structural Evolution of a Classical Archean Orogenic Lode-Gold System

Scott Anderson, Doug Berthelsen and Shawn Horte

Volume 39, Number 4, 2012

URI: https://id.erudit.org/iderudit/geocan39_4fs01

See table of contents

Publisher(s)

The Geological Association of Canada

ISSN

0315-0941 (print) 1911-4850 (digital)

Explore this journal

Cite this document

Anderson, S., Berthelsen, D. & Horte, S. (2012). The Rice Lake Mine Trend, Manitoba:: Regional Setting, Host-rock Stratigraphy, and Structural Evolution of a Classical Archean Orogenic Lode-Gold System. *Geoscience Canada*, 39(4), 174–175.

All rights reserved ${\rm @}$ The Geological Association of Canada, 2012

This document is protected by copyright law. Use of the services of Érudit (including reproduction) is subject to its terms and conditions, which can be viewed online.

https://apropos.erudit.org/en/users/policy-on-use/



GAC-MAC 2013:

FIELD GUIDE SUMMARY

The Rice Lake Mine Trend, Manitoba: Regional Setting, Host-rock Stratigraphy, and Structural Evolution of a Classical Archean Orogenic Lode-Gold System

GAC-MAC Winnipeg 2013, pre-meeting field trip

Scott Anderson¹, Doug Berthelsen² and Shawn Horte²

¹Manitoba Geological Survey 360–1395 Ellice Avenue Winnipeg, MB, Canada, R3G 3P2 E-mail: scott.anderson@gov.mb.ca

²San Gold Corporation P.O. Box 1000 Bissett, MB, Canada, R0E 0J0

FIELD TRIP OBJECTIVES

This three-day field trip will provide an overview of the key geological attributes of a major orogenic gold system in the Archean Rice Lake greenstone belt of the Uchi Domain, western Superior Province. Situated along the crustal-scale Wanipigow Fault, the Rice Lake belt is thought to represent vestiges of a long-lived oceanic basin (ca. 2.9 – 2.7 Ga) that accreted to the south margin of the ca. 3.0 Ga continental North Caribou Terrane during tectonic assembly of the Superior Province.

With a collective gold endowment of approximately 5 million ounces, the Rice Lake mine trend is the most significant lode-gold district in Manitoba. Clean bedrock exposures at Rice Lake (Fig. 1) provide an excellent opportunity to examine: 1) the character and stratigraphy of Neoarchean subaqueous effusive volcanic and vol-



Figure 1. Rice Lake with the town of Bissett and the Rice Lake mine complex in background.



Figure 2. Intermediate volcanic conglomerate, Townsite unit.

GEOSCIENCE CANADA Volume 39 2012



Figure 3. Polymictic conglomerate ("Timiskaming-type"), Wanipigow Fault.



Figure 4. Fault-fill breccia vein and peripheral extension veins, Rice Lake mine.

caniclastic rocks (Fig. 2), associated subvolcanic intrusions, and an unconformably overlying succession of fluvial-alluvial ('Timiskaming-type') sedimentary rocks (Fig. 3); 2) the geometry and overprinting relationships of mesoscopic deformation structures, which demonstrate a protracted history of synvolcanic faulting, early regional tilting, local synorogenic extension, intense regional compression and lateorogenic dextral transcurrent shear; 3) complex arrays of auriferous quartzcarbonate veins hosted by shear and tensile fractures (Fig. 4); and 4) the effects of hydrothermal alteration under greenschist facies metamorphic conditions in a variety of rock types. This field trip will also include a visit to the Rice Lake mine complex and an

underground tour of spectacular new exposures of high-grade quartz-carbonate vein systems.

The field trip is designed for specialist geoscientists and graduate students interested in greenstone belt geology and orogenic gold deposits. It will provide opportunities to discuss the setting and controls of vein emplacement within the context of the geodynamic and metallogenic evolution of greenstone belts in general and the Uchi Domain in particular. The field trip complements the GAC-MAC 2013 Special Session entitled Tectonic Evolution and Metallogeny of the Superior Province: New Insights from the North Caribou Terrane, the 'Ring of Fire', and Beyond.

OTHER INFORMATION

Rice Lake is approximately 250 km driving distance from Winnipeg (3 hours travel time), in southeastern Manitoba. The field trip stops are easily accessible along roads or bush trails, but may require walking on uneven or wet surfaces; one stop will require a round-trip hike of 3.0 km on flat terrain, which may be wet in places. Personal protective equipment will be provided for the underground tour.

About our field trip summaries:

Geoscience Canada intends to publish summaries of forthcoming field trip guides that are national or international in scope. Authors may wish to submit summaries of these guides in order to 'advertise' their trip to the broad geoscience community. They are strongly advised to do so well in advance of the field trip. In particular, we welcome summaries from field trips that accompany forthcoming GAC–MAC annual meetings, or divisional meetings. Here, we publish summaries from seven of the field trips to be held in conjunction with the GAC–MAC 2013 annual meeting to be held in Winnipeg May 22nd-24th.