Geoscience Canada

Journal of the Geological Association of Canada Journal de l'Association Géologique du Canada



Sedimentological, Structural and Paleontological Highlights of the Carboniferous Maritimes Basin in Southern New Brunswick

David Keighly, Adrian Park and John Waldron

Volume 40, Number 4, 2013

URI: https://id.erudit.org/iderudit/1021073ar

See table of contents

Publisher(s)

The Geological Association of Canada

ISSN

0315-0941 (print) 1911-4850 (digital)

Explore this journal

Cite this document

Keighly, D., Park, A. & Waldron, J. (2013). Sedimentological, Structural and Paleontological Highlights of the Carboniferous Maritimes Basin in Southern New Brunswick. *Geoscience Canada*, 40(4), 359–360.

All Rights Reserved © The Geological Association of Canada, 2013

This document is protected by copyright law. Use of the services of Érudit (including reproduction) is subject to its terms and conditions, which can be viewed online.

https://apropos.erudit.org/en/users/policy-on-use/



GEOSCIENCE CANADA Volume 40 2013

GAC-MAC 2014: FIELD GUIDE SUMMARY

Sedimentological,
Structural and
Paleontological Highlights
of the Carboniferous
Maritimes Basin in
Southern New Brunswick

GAC-MAC Fredericton 2014, post-meeting field trip

David Keighly¹, Adrian Park¹ and John Waldron²

'Department of Earth Science University of New Brunswick P.O. Box 4000 Fredericton, NB, Canada keig@unb.ca

²Department of Earth and Atmospheric Sciences University of Alberta Edmonton, Alberta, Canada, T6G 2E3

FIELD TRIP OBJECTIVES

Over three days, this excursion will offer participants the opportunity to visit spectacular Carboniferous outcrops of palaeontologic, sedimentologic and structural interest in southern New Brunswick. The Norton fossil forest (Fig. 1) contains upright tree trunks smaller, but considerably older than at Joggins. Near Sussex, soft-sediment deformation (slumping) of shale and oil shale is visible over several magnitudes, with these strata overlain by coarsening upward lake shoreface



Figure 1. Tournaisian upright trees, Norton fossil forest, New Brunswick.

sandstone. At Albert Mines, solid bitumen (Albertite) is abundant adjacent to abandoned mine shafts and fossil fish may be excavated with a bit of luck. Thick beds of often imbricate alluvial fan conglomerate form the 'flower

pots' at Hopewell Cape (Fig. 2). Salt diapirism (Fig. 3) has folded and faulted over a kilometre thickness of nonmarine strata on the Maringouin peninsula.




Figure 2. Flower pots at Hopewell Cape, New Brunswick.



Figure 3. Outcropping deformed gypsum associated with salt diapirism, Maringouin peninsula, New Brunswick.

ADDITIONAL INFORMATION

The trip will depart UNB on the Saturday morning of May 24th and returns to Fredericton on May 26th. Although this excursion will visit accessible roadside exposures, there will be significant walking along beaches (partly over boulders, and approximately 200 stairs), including one traverse 6 km (round trip) in length. Due to uneven terrain and unpredictable weather, rugged footwear and rain gear are highly recommended. Two night's accommodation in Moncton and meals will be included in the field trip registration fee.