

## Distribution of plant-parasitic nematodes in strawberry and raspberry fields in Quebec

G. Bélair et S. Khanizadeh

Volume 75, numéro 2, 1994

URI : <https://id.erudit.org/iderudit/706056ar>

DOI : <https://doi.org/10.7202/706056ar>

[Aller au sommaire du numéro](#)

Éditeur(s)

Société de protection des plantes du Québec (SPPQ)

ISSN

0031-9511 (imprimé)

1710-1603 (numérique)

[Découvrir la revue](#)

Citer cet article

Bélair, G. & Khanizadeh, S. (1994). Distribution of plant-parasitic nematodes in strawberry and raspberry fields in Quebec. *Phytoprotection*, 75(2), 101–107. <https://doi.org/10.7202/706056ar>

Résumé de l'article

Afin de connaître les nematodes phytoparasites associés au fraisier (*Fragaria ananassa*) et au framboisier (*Rubus idaeus*) dans sept régions agricoles du Québec (Cantons de l'Est, L'Assomption, Mauricie, Nicolet, Québec, Richelieu et Saint-Jean), 213 échantillons de sol ont été récoltés et analysés en juin et en juillet 1980. Le genre *Pratylenchus* fut le plus communément rencontré, soit dans 56,3% des champs de fraise et 46,4% des champs de framboise échantillonnés. Les densités de population du genre *Pratylenchus* étaient négativement corrélées avec les densités du *Meloidogynehapla* et le pH du sol. Le *Pratylenchus* était absent des champs de fraise des régions de Richelieu et de la Mauricie. Le *M. hapla* a été relevé dans des champs de fraise des régions de L'Assomption, Nicolet, Québec et Saint-Jean, et dans des champs de framboise des régions de Québec et de Saint-Jean. La présence du genre *Xiphinema* a été observée uniquement dans un champ de framboise des Cantons de l'Est. Les genres *Tylenchus*, *Tylenchorhynchus* et *Helicotylenchus* n'ont pas été observés dans la framboise. Les régions agricoles ont été divisées en groupes homogènes en fonction de leur distribution de nematodes, telle que révélée par une analyse en grappe.

---

## Distribution of plant-parasitic nematodes in strawberry and raspberry fields in Quebec

Guy Bélair<sup>1</sup> and Shahrokh Khanizadeh<sup>1,2</sup>

Received 1993-09-14; accepted 1994-11-01

---

In a survey of plant-parasitic nematodes associated with strawberry (*Fragaria x ananassa*) and raspberry (*Rubus idaeus*) in seven Quebec agricultural regions (Eastern Townships, L'Assomption, Mauricie, Nicolet, Québec, Richelieu, and Saint-Jean), 213 soil samples were analyzed in June and July 1980. The most commonly encountered nematode genus was *Pratylenchus*, occurring in 56.3 % and 46.4% of sampled strawberry and raspberry fields, respectively. The population density of *Pratylenchus* was negatively correlated with that of *Meloidogyne hapla* and with soil pH. No *Pratylenchus* was found in raspberry fields from the Mauricie and Richelieu regions. *M. hapla* was found in strawberry fields of the L'Assomption, Nicolet, Québec and Saint-Jean regions, and also in raspberry fields in the Québec and Saint-Jean regions. The genus *Xiphinema* was recorded in only one raspberry field in the Eastern Townships. *Tylenchus*, *Tylenchorhynchus* and *Helicotylenchus* were not found on raspberry. Agricultural regions were divided into homogeneous groups with respect to their nematode distribution as revealed by cluster analysis.

**Bélair, G. et S. Khanizadeh. 1994. Distribution des nématodes phytoparasites dans les champs de fraise et de framboise au Québec. PHYTOPROTECTION 75: 101-107.**

Afin de connaître les nématodes phytoparasites associés au fraisier (*Fragaria x ananassa*) et au framboisier (*Rubus idaeus*) dans sept régions agricoles du Québec (Cantons de l'Est, L'Assomption, Mauricie, Nicolet, Québec, Richelieu et Saint-Jean), 213 échantillons de sol ont été récoltés et analysés en juin et en juillet 1980. Le genre *Pratylenchus* fut le plus communément rencontré, soit dans 56,3% des champs de fraise et 46,4% des champs de framboise échantillonnés. Les densités de population du genre *Pratylenchus* étaient négativement corrélées avec les densités du *Meloidogyne hapla* et le pH du sol. Le *Pratylenchus* était absent des champs de fraise des régions de Richelieu et de la Mauricie. Le *M. hapla* a été relevé dans des champs de fraise des régions de L'Assomption, Nicolet, Québec et Saint-Jean, et dans des champs de framboise des régions de Québec et de Saint-Jean. La présence du genre *Xiphinema* a été observée uniquement dans un champ de framboise des Cantons de l'Est. Les genres *Tylenchus*, *Tylenchorhynchus* et *Helicotylenchus* n'ont pas été observés dans la framboise. Les régions agricoles ont été divisées en groupes homogènes en fonction de leur distribution de nématodes, telle que révélée par une analyse en grappe.

1. Agriculture and Agri-Food Canada, Horticulture research and Development Centre, 430, Gouin blvd., Saint-Jean-sur-Richelieu, Quebec, Canada J3B 3E6. Contribution N° 335/94.10.02R
2. Department of Plant Science, Macdonald Campus of McGill University, Sainte-Anne-de-Bellevue, Quebec, Canada H9X 3V9

## INTRODUCTION

Quebec is a major berry-producing area in eastern Canada with 11 263 t of strawberry (*Fragaria x ananassa* Duch.) and 931 t of raspberry (*Rubus idaeus* L.) for a \$26 million crop return annually (Statistics Canada 1992). These two small fruit crops are produced on a wide range of soil types with different cultural practices. While some growers are unaware of nematode problems, others use nematicides on a routine basis. Root-lesion nematodes (*Pratylenchus* spp.) and root-knot nematodes (*Meloidogyne hapla* Chitwood) have been isolated from strawberry and raspberry fields from soil samples processed through the Nematology Advisory Service (Agriculture and Agri-Food Canada). However, extensive surveys had not been conducted to date in Quebec berry-producing areas.

Surveys conducted from 1972-1979 in the province have identified the major plant-parasitic nematodes on apple (*Malus pumila* Borkh.), tobacco (*Nicotiana tabacum* L.), forage and vegetable crops (Meloche *et al.* 1980; Santerre and Lévesque 1982; Vrain and Dupré 1982; Vrain and Rousselle 1980; Willis *et al.* 1976). In most agricultural regions, the root-lesion nematode, *Pratylenchus* spp., was the most dominant genus, occasionally reaching damaging population densities.

This survey was conducted to determine the density and distribution of plant-parasitic nematodes in the major berry-producing areas of Quebec; to assess any possible correlation between the densities of these organisms and specific abiotic variables recorded in this survey; and to determine any similarities between agricultural regions in terms of nematode types and densities.

## MATERIALS AND METHODS

During June and July 1980, a total of 213 soil samples were recovered from 185 strawberry and 28 raspberry fields in a survey covering the seven major berry-producing areas in Quebec (Eastern Townships, L'Assomption,

Mauricie, Nicolet, Québec, Richelieu, and Saint-Jean). Although fields were not selected using strict random procedure, no effort was made to select fields with conspicuous problems. Ten to fifteen soil subsamples (0-20 cm deep) taken at random in the row with a hand trowel were combined to form a composite sample for each field of 0.2-0.8 ha. Soil samples were put into plastic bags and stored at 10-15°C until needed.

Nematode soil densities were determined by extracting nematodes from a 100-cm<sup>3</sup> subsample using the modified Baermann funnel method (Townshend 1963). Identification to genus and nematode counts were made using a stereomicroscope. Specific identifications were verified by Dr. R.V. Anderson, Biosystematics Research Center, Ottawa, Ontario. Nematode counts were transformed to log (x + 2) prior to analysis as proposed by Proctor and Marks (1974) and soil samples were analyzed for sand fraction (Bouyoucos 1962). Soil pH and organic content were determined by using standard procedures (McKeague 1977).

Simple statistics (mean, maximum, and frequency) and cluster analysis were performed on nematode counts using SAS (SAS Institute Inc. 1988). The cluster analysis was used to distinguish clusters of regions that have similar attributes with respect to nematode genera and density. Arcsine transformation of percentages were used prior to analysis when needed.

## RESULTS

Eight genera of stylet-bearing nematodes were isolated from strawberry soils and six genera were isolated from raspberry soils in Quebec (Tables 1, 2). The root-lesion nematodes *Pratylenchus penetrans* Cobb and *P. crenatus* Loof were the most conspicuous plant-parasitic nematodes associated with these crops, and were present in 56.3 % and 46.4% of sampled strawberry and raspberry fields, respectively. Of the soil samples infested with *Pratylenchus* spp., 94.0% had *P. penetrans*, 7.7% had *P. crenatus*, and only 1.7% had both species.

The northern root-knot nematode, *Meloidogyne hapla*, was found in 11.9% and 7.1% of the sampled strawberry and raspberry fields, respectively. When present, *M. hapla* was found in greater population densities in raspberry than in strawberry. *M. hapla* was not found in soil samples from the Eastern Town-

ships, Mauricie and Richelieu regions, neither in raspberry fields in L'Assomption and Nicolet. The pin nematode, *Paratylenchus* spp., was the most numerous plant-parasitic nematode in all tested strawberry fields. *Paratylenchus* spp. was not found in raspberry fields in the Mauricie and Richelieu

**Table 1. Number of fields, mean and maximum density of plant-parasitic nematodes, and frequency of infestation in 185 strawberry fields in seven agricultural regions of the Quebec province**

Agricultural region	Eastern Townships	L'Assomption	Mauricie	Nicolet	Québec	Richelieu	Saint-Jean	Overall mean and total frequency	Total infested fields (%)
N <sup>a</sup>	13	57	10	12	65	9	19		
<i>Pratylenchus</i> spp.									
Mean <sup>b</sup>	53	26	46	52	38	5	13	33	
Maximum <sup>b</sup>	180	300	120	290	238	10	28		
Frequency <sup>c</sup>	11	27	8	8	43	4	3	104	56.3
<i>Meloidogyne</i> spp.									
Mean	0	10	0	28	7	0	3	7	
Maximum	0	22	0	28	14	0	4		
Frequency	0	10	0	1	9	0	2	22	11.9
<i>Paratylenchus</i> spp.									
Mean	93	11	25	176	8	4	7	46	
Maximum	350	30	70	590	16	4	12		
Frequency	4	8	4	4	6	1	4	31	16.8
<i>Tylenchus</i> spp.									
Mean	0	5	6	0	34	0	0	6	
Maximum	0	5	6	0	120	0	0		
Frequency	0	2	1	0	4	0	0	7	3.8
<i>Trichodorus</i> spp.									
Mean <sup>b</sup>	0	6	0	0	7	15	0	4	
Maximum	0	8	0	0	12	26	0		
Frequency	0	3	0	0	3	2	0	8	4.3
<i>Tylenchorhynchus</i> spp.									
Mean	4	11	0	0	0	0	0	2	
Maximum	43	16	0	0	0	0	0		
Frequency	1	2	0	0	0	0	0	3	1.6
<i>Helicotylenchus</i> spp.									
Mean	0	0	0	0	4	0	0	1	
Maximum	0	0	0	0	4	0	0	4	
Frequency	0	0	0	0	1	0	0	1	0.5
<i>Ditylenchus</i> spp.									
Mean	0	4	4	0	3	4	2	2	
Maximum	0	4	4	0	3	4	2	17	
Frequency	0	2	1	0	1	1	1	6	3.2

<sup>a</sup> Number of fields sampled in each agricultural region.

<sup>b</sup> Number of nematodes in 100 cm<sup>3</sup> of soil.

<sup>c</sup> Number of infested fields.

**Table 2. Number of fields, mean and maximum density of plant-parasitic nematodes, and frequency of infestation in 28 raspberry fields in seven agricultural regions of the Quebec province**

Agricultural region	Eastern Townships	L'Assomption	Mauricie	Nicolet	Québec	Richelieu	Saint-Jean	Overall mean and total frequency	Total infested fields (%)
N <sup>a</sup>	4	12	1	1	3	2	5		
<i>Pratylenchus</i> spp.									
Mean <sup>b</sup>	45	12	12	5	12	26	34	21	
Maximum <sup>b</sup>	88	20	12	5	12	32	40		
Frequency <sup>c</sup>	3	3	1	1	1	2	2	13	46.4
<i>Meloidogyne</i> spp.									
Mean	0	0	0	0	48	0	42	13	
Maximum	0	0	0	0	48	0	42		
Frequency	0	0	0	0	1	0	1	2	7.1
<i>Paratylenchus</i> spp.									
Mean	22	69	0	4	200	0	18	45	
Maximum	35	138	0	4	200	0	21		
Frequency	2	3	0	1	1	0	2	9	32.1
<i>Trichodorus</i> spp.									
Mean	2	0	0	0	0	0	0	0.3	
Maximum	2	0	0	0	0	0	0		
Frequency	1	0	0	0	0	0	0	1	3.6
<i>Ditylenchus</i> spp.									
Mean	15	0	0	0	0	0	0	2	
Maximum	15	0	0	0	0	0	0		
Frequency	1	0	0	0	0	0	0	1	3.6
<i>Xiphinema</i> spp.									
Mean	250	0	0	0	0	0	0	36	
Maximum	250	0	0	0	0	0	0		
Frequency	1	0	0	0	0	0	0	1	3.6

<sup>a</sup> Number of fields sampled in each agricultural region.

<sup>b</sup> Number of nematodes in 100 cm<sup>3</sup> of soil.

<sup>c</sup> Number of infested fields.

regions. No information is presently available on the pathogenicity of this ectoparasitic genus on raspberry and strawberry plants. No *Tylenchus* spp., *Tylenchorhynchus* spp. and *Helicotylenchus* spp. were observed in raspberry soil samples collected. *Trichodorus* spp. and *Ditylenchus* spp. were found in a raspberry field in the Eastern Townships. *Xiphinema rivesi* Dalmasso was confined to only one raspberry field in the Eastern Townships and was not observed in any strawberry fields. *Tylenchus* spp. was associated with strawberry fields in the L'Assomption, Mauricie and Québec regions (Table 1).

*Trichodorus* spp. was found only in strawberry fields in the L'Assomption, Québec and Richelieu regions. *Tylenchorhynchus* spp. was observed in the Eastern townships and L'Assomption, and *Helicotylenchus* spp. was observed in strawberry fields in the Québec region. *Ditylenchus* spp. was not observed in strawberry fields in the Eastern Townships and Nicolet regions.

Significant correlations were detected among abiotic variables and nematode densities. The population density of *Pratylenchus* spp. was negatively correlated with *M. hapla* ( $r = -0.26$ ;

$P < 0.01$ ) and with soil pH ( $r = -0.23$ ;  $P < 0.01$ ) but was positively correlated ( $r = 0.19$ ;  $P < 0.05$ ) with *Paratylenchus* spp. The number of *Paratylenchus* spp. was positively correlated ( $r = 0.15$ ;  $P < 0.05$ ) with the organic matter content. No significant correlation was established between plant-parasitic nematodes and sand content ( $r = -0.06$ ;  $P > 0.05$ ).

Cluster analysis revealed that the seven regions can be divided into two homogeneous groups with respect to the nematode distribution in strawberry fields. Group 1 consists of the Eastern Townships, Mauricie, and Nicolet regions and group 2 consists of the L'Assomption, Richelieu, Saint-Jean, and Québec regions (Fig. 1A). A similar procedure was applied to the data collected from raspberry fields. The seven agricultural regions were also divided into two groups: group 1 comprising Québec and the Eastern Townships and group 2 comprising all other regions (Fig. 1B).

## DISCUSSION

*Pratylenchus* spp. is by far the most dominant plant-parasitic nematode both on strawberry and raspberry in Québec. *P. penetrans* and *P. crenatus* were found in association but *P. penetrans* prevailed. This widespread distribution of *Pratylenchus* spp. in all agricultural regions is consistent with results from earlier surveys of apple orchard and vegetable crops in Québec (Vrain and Dupré 1982; Vrain and Rousselle 1980). Nematicide treatments provided evidence that *P. penetrans* populations can impair growth and yield of raspberry and strawberry in Eastern Canada (Bélair 1991; Kimpinski 1985).

The northern root-knot nematode, *M. hapla*, also shown to be pathogenic to both strawberry and raspberry (McElroy 1972), is present on both crops in four agricultural regions. A similar proportion of *M. hapla* infested soils was found in vegetable fields of southern Québec (Vrain and Dupré 1982). Because *M. hapla* soil population densities drop annually below the detection level in

July (G. Bélair, unpublished data), the actual distribution of this nematode may have been underestimated in this survey.

The dagger nematode, *Xiphinema* spp., was not isolated from strawberry fields but was isolated from one raspberry field from the Eastern Townships region where small fruits are occasionally grown on apple orchard soils. A nematological survey of Québec orchards revealed the widespread distribution of *X. rivesi* in all apple-producing areas. In Québec, there is some indication that *X. rivesi* could be associated with raspberry decline (Bélair 1991).

Correlations between nematode and abiotic variables were rarely observed. Weak correlations could be due to large variations in nematode counts resulting from soil population changes that occurred during the sampling period.

This survey identified *Pratylenchus* spp. and *M. hapla* as the most important nematode parasites in Québec strawberry and raspberry plantations. Most raspberry fields in all agricultural regions had similar nematode populations. However, strawberry fields in the L'Assomption, Richelieu, Saint-Jean and Québec regions had different nematode populations compared to the Eastern Townships, Mauricie and Nicolet regions. Similarities in the distribution of plant-parasitic nematodes among regions suggests that factors such as cultural practices, including crop rotation, cultivars, and weed management could be involved. The homogeneous group of strawberry fields made of the L'Assomption, Richelieu, Saint-Jean and Québec regions was characterized by a greater diversity of the plant-parasitic nematodes. In these four agricultural regions, growers include various vegetable crops such as corn (*Zea mays* L.) and potato (*Solanum tuberosum* L.) in their rotation program, rather than small grains or buckwheat (*Fagopyrum esculentum* L.) which are predominantly used in the three other producing areas.

Because cultural practices within each agricultural region have remained basically unchanged for the last 10 yr, we




Figure 1. Ward minimum variance cluster analysis based on genera and nematode number collected from (A) strawberry and (B) raspberry fields of seven agricultural regions in Quebec.

believe that this survey provides a general and realistic representation of the nematode distribution in small fruit production areas of the province of Quebec. The redistribution of some plant-parasitic nematodes from contaminated transplants remains possible since the Quebec certification program does not take nematodes into account.

The importance of these nematodes in strawberry and raspberry fields has not been evaluated yet. More intensive samplings and yield estimates are required to have a better understanding of the relative importance of the plant-parasitic nematodes in small fruit production.

## REFERENCES

- Bélaïr, G. 1991.** Effects of preplant soil fumigation on nematode population densities and on growth and yield of raspberry. *Phytoprotection* 72: 21-25.
- Bouyoucos, G.J. 1962.** Hydrometer method improved for making particle size analysis of soils. *Agron. J.* 54: 464-465.
- Kimpinski, J. 1985.** Nematodes in strawberries on Prince Edward Island, Canada. *Plant Dis.* 69: 105-107.
- McElroy, F.D. 1972.** Nematodes of tree fruits and small fruits. Pages 335-376 in J.M. Webster (ed.), *Economic nematology*. Acad. Press, New York.
- McKeague, J.A. 1977.** Manuel de méthodes d'échantillonnage et d'analyse des sols. Comité canadien de pédologie, Soil Research Institute, Agriculture Canada, Ottawa. 250 pp.
- Meloche, F., J.-G. Pilon, G. Mailloux, and T.C. Vrain. 1980.** Inventaire des problèmes entomologiques et nématologiques dans les plantations de tabac jaune au Québec. *Ann. Soc. Entomol. Que.* 25: 81-89.
- Proctor, J.R., and C.F. Marks. 1974.** The determination of normalizing transformations for nematode count data from soil samples and of efficient sampling schemes. *Nematologica* 20: 395-406.
- Santerre, J., and R. Lévesque. 1982.** Inventaire de nématodes phytoparasites dans des cultures de plantes fourragères au Québec: 1973 à 1978. *Can. Plant Dis. Surv.* 62: 13-19.
- SAS Institute Inc. 1988.** SAS/STAT User's guide: release 6.03. SAS Institute Inc. Cary, North Carolina. 1029 pp.
- Statistics Canada. 1992.** Fruit and vegetable production. Ministry of supply and services Canada 1992, catalogue 22-003, vol. 61, n° 2, 32 pp.
- Townshend, J.L. 1963.** A modification and evaluation of the apparatus for the Oostenbrink direct cottonwool filter extraction method. *Nematologica* 9: 106-110.
- Vrain, T.C., and M. Dupré. 1982.** Distribution des nématodes phytoparasites dans les sols maraîchers du sud-ouest du Québec. *Phytoprotection* 63: 79-85.
- Vrain, T.C., and G.L. Rousselle. 1980.** Distribution of plant-parasitic nematodes in Quebec apple orchards. *Plant Dis.* 64: 582-583.
- Willis, C.B., J.L. Townshend, R.V. Anderson, J. Kimpinski, R.H. Mulvey, J.W. Potter, J. Santerre, and L.Y. Wu. 1976.** Species of plant-parasitic nematodes associated with forage crops in Eastern Canada. *Plant Dis. Rep.* 60: 207-210.